

Rare Plants Found In and Near Canfor's FMA Area
(Provincial and Global ranking from ANHIC 1999)

Scientific Name	Common Name	SRank	GRank	Discovered in:
<i>Agrostis exarata</i>	spike redtop	S2	G5	5
<i>Agrostis humilis</i>	low bent grass	S1S2	G4	1
<i>Alium gezeri</i>	Geyer's onion	S2	G4G5	4
<i>Aquilegia Formosa</i>	Sitka columbine	S2	G5	4
<i>Arnica amplexicaulis</i>	stem-clasping arnica	S2	G4	4
<i>Artemisia furcata</i> var. <i>furcata</i>	forked wormweed	S1	G4T?	4
<i>Aster x maccallae</i>	n/a	S1S2	HYB	4
<i>Atrichum undulatum</i>	undulated crane's bill moss	S1S2	G5	1
<i>Botrychium multifidum</i> var. <i>intermedium</i>	leather grape fern	S2	G5T4?	5
<i>Botrychium simplex</i>	dwarf grape fern	S1S2	G5	5
<i>Brachythecium campestre</i>	n/a	S2	G4G5Q	1
<i>Brachythecium hylotapetum</i>	n/a	S2	G?	1
<i>Brachythecium leibergii</i>	n/a	S2	G?	1
<i>Bryum pallens</i>	n/a	S2	G4G5	2
<i>Calamagrostis lapponica</i>	Lapland reed grass	S1	G5	4
<i>Carex mertensii</i>	purple sedge	S1	G5	2,5
<i>Carex retrorsa</i>	turned sedge	S2	G5	2
<i>Carex rostrata</i>	beaked sedge	S2	G5	1,3,4
<i>Cypripedium acaule</i>	stemless lady's-slipper	S2	G5	1
<i>Cystopteris montana</i>	mountain bladder fern	S2	G5	5
<i>Drosera anglica</i>	oblong-leaved sundew	S2	G5	1
<i>Epilobium glaberrimum</i> ssp. <i>fastigiatum</i>	willowherb	S1	G5T?	1,3,4
<i>Fontinalis antipyretica</i>	n/a	S2	G5	2
<i>Gentiana glauca</i>	alpine gentian	S2	G4G5	4
<i>Hieracium cynoglossoides</i>	wolly hawkweed	S2	G5?	1,4
<i>Lactuca biennis</i>	tall blue lettuce	S2S3	G5	2
<i>Luzula acuminata</i>	wood-rush	S1	G5	5
<i>Luzula rufescens</i>	reddish wood-rush	S1	G5	5
<i>Lycopodium selago</i>	mountain club-moss	S1	G5	1,3
<i>Lycopodium stichense</i>	ground-fir	S2	G5	5
<i>Malaxis monophylla</i>	white adder's-mouth	S2	G5	1,3
<i>Mimulus guttatus</i>	yellow monkeyflower	SU	G5	1
<i>Orthotrichum pallens</i>	n/a	S2	G5	2
<i>Oryzopsis exigua</i>	little rice grass	S1	G5	4
<i>Osmorhiza longistylis</i>	smooth sweet cicely	S2	G5	4
<i>Osmorhiza purpurea</i>	purple sweet cicely	S2	G4G5	1,4
<i>Pedicularis capitata</i>	large-flowered lousewort	S2	G4	4
<i>Phegopteris connectilis</i>	northern beech fern	S2	G5	2,5
<i>Potamogeton foliosus</i>	leafy pond weed	S2	G5	2
<i>Potentilla drummondii</i>	Drummond's cinquefoil	S2	G5	5
<i>Potentilla hookeriana</i>	Hooker's cinquefoil	S2	G4	4
<i>Potentilla multifida</i>	branched cinquefoil	S1	G5	4
<i>Pyrola grandiflora</i>	Arctic wintergreen	S2	G5	1,4
<i>Pyrola picta</i>	white-veined wintergreen	S1	G4G5	4
<i>Ranunculus occidentalis</i> var. <i>brevistylis</i>	western butter cup	S2	G5T5	5
<i>Rhizomnium nudum</i>	n/a	S2	G?	1
<i>Rhytiadelphus squarrosus</i>	pipecleaner moss	S2	G4G5	1
<i>Salix alaxensis</i> var. <i>alaxensis</i>	Alaska willow	S2	G5T?	4
<i>Salix commutata</i>	changeable willow	S2	G5	1,4
<i>Salix raupii</i>	Raup's willow	S1	G2	5
<i>Salix stonlonifera</i>	willow	S1	G4G5	4
<i>Scouleria aquatica</i>	n/a	S2	G4	1
<i>Sphenopholis obtusata</i>	prairie wedge grass	S2	G5	2
<i>Stellaria crispa</i>	wavy-leaved chickweed	S2	G5	2,5
<i>Streptopus roseus</i>	rose mandarin	S1	G5	1,2,3,4
<i>Streptopus streptopoides</i>	twisted-stalk	S1	G5	5
<i>Thuidium delicatulum</i>	n/a	SR	G5	1
<i>Townsendia exscapa</i>	low townsendia	S1S2	G5	5
<i>Vaccinium uliginosum</i>	bog bilberry	S2	G5	5

Rank: S = Subnational level (status in a province, i.e., Alberta) G = Global level (status throughout its global range)

S1/G1 - Critically imperiled due to extreme rarity (5 or fewer occurrences), S2/G2 - Imperiled because of rarity (6 to 20 occurrences), S3/G3 - Rare or uncommon (21 to 100 occurrences), S4/G4 - Apparently secure, with many occurrences, S5/G5 - Abundant and secure, with many occurrences, SR - Reported but without persuasive documentation to either accept or reject the report, SU - Uncertain status, possibly in peril; more information is required, T - Rank for subspecific taxon (subspecies or variety), Q - Taxonomic problems involved; more information is required, HYB - Hybrid taxon that is recurrent in the landscape, ? - Rank tentatively assigned; no information is available or the number of occurrences is estimated.

Discovered in: 1. 1643 (AEP and Canfor) plots sampled inside and within 20 km buffer of FMA area, 2. a query of the ANHIC database for the requested area (townships listed in Appendix ?), 3. 1997 Canfor plots only (according to Snyder 1998), 4. AEP plots in the region (according to Snyder 1998), 5. Snyder (1998): species not found in either AEP or Canfor plots but listed in the region