

Salinity Mapping for Resource Management

within the M.D. of Rocky View, Alberta

J. Kwiatkowski
L.C. Marciak
D. Wentz
C.R. King

Conservation and Development Branch
Alberta Agriculture, Food and Rural Development

March 1996

Canada


Agriculture - Building a Healthy Environment

Alberta

Abstract

This report presents a methodology to map salinity at a municipal scale and applies this procedure to the M.D. of Rocky View, a municipality in southern Alberta. The methodology was developed for the County of Vulcan (Kwiatkowski et al. 1994) and is being applied to other Alberta municipalities which have identified soil salinity as a concern.

Soil salinity is a major land degradation issue in the M.D. of Rocky View. The information on salinity location, extent, type and control measures presented in this report will help the municipality's planners to target salinity control and resource management programs.

The methodology has four steps:

1. The location and extent of saline areas are mapped based on existing information including aerial photographs, maps, satellite imagery and information from local personnel and field inspections.
2. Saline areas are classified on the basis of the mechanism causing salinity. The mechanism is important because it determines which control measures are appropriate. Eight salinity types are recognized within Alberta. These are: contact/slope change salinity, outcrop salinity, artesian salinity, depression bottom salinity, coulee bottom salinity, slough ring, irrigation canal seepage salinity and natural/irrigation salinity. All eight types occur in the M.D. of Rocky View.
3. Cost-effective, practical control measures are identified for each salinity type.
4. Colour-coded maps at 1:100 000 and 1:200 000 are prepared showing salinity location, extent and type. Also, a digital data base is compiled in ArcInfo format. This data base can be easily managed and transformed into ArcView format.

Analysis of the mapping data shows that 915 saline areas occur in the M.D. and these areas occupy a total of 11 358 ha (28 068 ac). This represents 2.5% of the M.D.'s total area. Only saline areas visible on the soil surface are mapped. This provides information needed by the municipality's staff to target sites for further investigation. The surrounding lands may have weakly saline subsurface soils which can reduce yields of sensitive crops. Thus salinity control practices may benefit crop yields over a much broader area than just the visible seep.

Depression bottom salinity is the most common salinity type (58.1% of saline land in the M.D.), followed by contact/slope change salinity (22.3% of saline land), irrigation canal seepage salinity (15.7%), irrigation/natural salinity (2.1%), coulee bottom salinity (0.9%), outcrop salinity (0.6%), slough ring salinity (0.2%), and artesian salinity (0.1%). There are many small contact/slope change seeps (554 seeps), and fewer but larger depression bottom and coulee bottom seeps (263 seeps and 8 seeps, respectively).

Acknowledgements

This salinity mapping and analysis project for the M.D. of Rocky View was managed by the Conservation and Development Branch of Alberta Agriculture, Food and Rural Development. Funding provided by the Conservation and Development Branch, the Canada-Alberta Environmentally Sustainable Agriculture (CAESA) Agreement and the M.D. of Rocky View is greatly appreciated.

The authors acknowledge the contributions made by: James Pittman and Eugene Kurinka, Conservation and Development Branch, Alberta Agriculture, Food and Development, for field work, aerial photograph interpretation and water sampling in the M.D. Rocky View; and Gerald Stark, Longin Pawlowski and Jason Miller, Conservation and Development Branch, Alberta Agriculture, Food and Rural Development, for preparing diagrams and logos, and plotting the salinity map. Comments on the salinity data from Tim Dietzler, Agricultural Fieldman, M.D. of Rocky View, are greatly appreciated.

Contents

	Page
Abstract	i
Acknowledgements	ii
Tables	iv
Figures	iv
1.0 Introduction	1
1.1 Goal and Objectives	1
1.2 Methodology	1
1.3 Information Sources	1
2.0 Classification and Management of Saline Seeps	3
2.1 Transportation of Salts	3
2.2 Salinity Types	4
2.2.1 Dryland Salinity Types	4
2.2.2 Irrigation Salinity Types	7
2.3 Salinity Controls	8
2.3.1 Biological Controls	8
2.3.2 Structural Controls	11
3.0 Saline Water Bodies	13
4.0 Saline Seep Distribution	15
5.0 References	20
Glossary	22

Tables

	Page
Table 1. Salinity types and control methods	9
Table 2. Forage crops for saline soils and flooded areas	10
Table 3. Salinity distribution by type in the M.D. of Rocky View	15

Figures

	Page
Figure 1. Generalized saline seep mechanisms	3
Figure 2. Contact salinity	4
Figure 3. Slope change salinity	5
Figure 4. Outcrop salinity	5
Figure 5. Artesian salinity	6
Figure 6. Depression bottom salinity	6
Figure 7. Slough ring salinity	7
Figure 8. Canal seepage salinity	8
Figure 9. Saline water bodies and associated salinity types, Twp. 27 Rge. 25 W4, M.D. of Rocky View	14
Figure 10. Number of saline seeps by type in the M.D. of Rocky View	17
Figure 11. Number and size of saline seeps in the M.D. of Rocky View	18
Figure 12. Number and size of contact/slope change saline seeps in the M.D. of Rocky View	19

1.0 Introduction

1.1 Goal and Objectives

The goal of this project is to present information on salinity location, extent and type in the Municipal District of Rocky View. Soil degradation is a key issue in conservation planning and salinity is one of the most visible soil degradation problems affecting the M.D. according to its Canada-Alberta Environmentally Sustainable Agriculture (CAESA) Agreement action plan. Thus this mapping project aims to better define the salinity problem for the M.D. of Rocky View.

The project's goal is achieved through the following objectives:

1. To derive and integrate existing salinity information for agricultural land in the M.D. of Rocky View.
2. To determine the salinity type based on the salinity mechanism.
3. To recommend appropriate control methods for each type of salinity.
4. To compile a map depicting salinity location, extent and type.

The project differs from most salinity surveys by specifying the type, exact locations and control measures. This information can be used in municipal and farm conservation planning.

1.2 Methodology

The methodology for mapping salinity was developed for the County of Vulcan by Kwiatkowski et al. (1994). It is being applied to other Alberta municipalities where soil salinity is a concern. The process of salinity mapping consists of four stages:

1. Scan aerial photographs and digitize saline areas on the municipal base map.
2. Determine the types of salinity occurring in the municipality, based on hydrogeology, surface water flow, geology, topography, irrigation and soils. Determine appropriate cost-effective, practical control measures based on salinity types.
3. Field check the salinity data and submit the draft salinity information to a technical team consisting of a project manager, hydrogeologist, salinity specialist, and the local Agricultural Fieldman and District Specialist for review.
4. Prepare a colour-coded 1:100 000 map, showing salinity location, extent and type, and an accompanying report with a 1:200 000 map. Also, a digital data base is compiled in ArcInfo format. This data base can be easily managed and transformed into ArcView format.

1.3 Information Sources

A variety of maps, aerial photographs and other information sources were used for this project. Information on climate, soils, parent material and geology is from the following soil survey and hydrogeology reports:

- Soil Survey of Rosebud and Banff Sheets (Wyatt et al. 1943)
- Soil Survey of Blackfoot and Calgary Sheets (Wyatt et al. 1960)
- Hydrogeology of the Calgary-Golden area, Alberta (Ozoray and Barnes 1978)
- Hydrogeology of the Gleichen area, Alberta (Ozoray and Lytviak 1974)
- Hydrogeology of the Drumheller area, Alberta (Borneuf 1972)

The Saline/Waterlogged Lands Map for the Western Irrigation District (1990) provided information on salinity in two categories: moderately and severely affected areas. The data were collected in 1982, 1985 and 1987 and include sloughs, stock-watering ponds and small, temporary water bodies. The main causes of salinization and waterlogging within irrigation districts are seepage from canals, poor water management, poor irrigation practices and inadequate drainage.

Aerial photographs from 1991 (scale of 1:30 000) were used to help determine the location, extent and type of salinity on a section-by-section basis. As well, satellite imagery was used, particularly to identify irrigated lands.

Prairie Farm Rehabilitation Administration (PFRA) Salinity Reports for 18 sites throughout the M.D. were also valuable information sources (Prairie Farm Rehabilitation Administration various dates). They provided detailed data including information on drilling investigations and severity and extent of salinity.

To ensure the accuracy of the salinity map, more than two-thirds of the municipality was field checked. Local personnel were also consulted to verify the findings.

2.0 Classification and Management of Saline Seeps

2.1 Transportation of Salts

The dominant salts in the M.D. of Rocky View consist of sodium and magnesium sulphates. Analyses of groundwater, saline soils and parent material suggest that the primary source of salts is bedrock, and the secondary source is glacial till (Greenlee et al. 1968). Soils developed on the Bearpaw bedrock formation contain high salt concentrations.

Saline seeps form when saline groundwater rises to the ground surface. Contact and slope change seeps (described in Section 2.2.1) develop when water in a recharge area percolates down through the soil profile beyond the root zone and dissolves soluble salts (Figure 1). The water moves laterally to a lower position in the landscape and through capillarity rises to the surface, resulting in a saline seep. High evapotranspiration rates cause the capillary rise and the deposition of salts on the soil surface.




Figure 1. Generalized saline seep mechanisms

Three different types of flow may be recognized within a groundwater basin: local, intermediate and regional. A local flow system occupies a relatively small area, with the recharge area at a higher elevation than the discharge area. An intermediate system consists of several interconnected local systems. A regional system has its recharge area at the water divide of a basin while the discharge area lies at the bottom of the basin. In the M.D. of Rocky View, most of the groundwater flow systems are local, and the recharge areas are within a few hundred metres of their discharge area. Intermediate flow systems extend beyond 1 km (0.6 mi) of their recharge area. Regional flow systems extend over several kilometres.

Groundwater movement is influenced by topography as follows:

- In large, flat areas, groundwater movement is minimal or even impeded.
- In areas with well-defined local relief (e.g. hummocky or rolling landscapes), local systems are prevalent.
- In areas with one large slope, regional systems are prevalent.
- In large valleys, regional systems predominate.

2.2 Salinity Types

Eight salinity types have been identified in Alberta (Kwiatkowski et al. 1994). They can be grouped into six dryland and two irrigation types. Based on hydrogeology, surface water flow, geology, topography, irrigation and soils, all eight types are found in the M.D. of Rocky View. Almost all saline seeps in the M.D. occur east of Highway 2.

2.2.1 Dryland Salinity Types

1. Contact/Slope Change Salinity

Contact salinity and slope change salinity are grouped together because they cannot be differentiated on aerial photographs and because the same methods are used to control both types (see Section 2.3). The two types are described as follows:

- a. **Contact salinity** occurs where a permeable water-bearing surface layer thins out above a less permeable layer (such as a fine textured layer). This forces the groundwater flow closer to the surface (Figure 2). Contact salinity is the second most common type of salinity in the M.D., occupying 22.3% of all saline lands. It dominates in sandy, gently rolling areas. It occurs throughout the eastern portion of the M.D. and is especially common east and southeast of Airdrie. Contact seeps also occur at the shoulders of coulees and as small seeps scattered throughout the M.D.




Figure 2. Contact salinity

- b. **Slope change salinity** occurs where the slope decreases. This decrease results in a slowing of the groundwater flow and a shallower water table (Figure 3). This type of seep expands upslope. It occurs in association with contact salinity type and is found throughout the eastern portion of the M.D. of Rocky View as a minor salinity type.




Figure 3. Slope change salinity

2. Outcrop Salinity

Outcrop salinity occurs where a permeable, water-bearing layer, such as a coal seam or fractured bedrock layer, outcrops at or near the surface (Figure 4). It is found along a slope at similar elevations. In the M.D. of Rocky View, this salinity type occupies only 0.6% of the total saline land. It is found in Twp. 26 Rge. 28, 29 W4, and in Twp. 26 Rge. 1 W5. It occurs mainly in conjunction with contact/slope change salinity.




Figure 4. Outcrop salinity

3. Artesian Salinity

Artesian salinity occurs where water from a pressurized aquifer rises to or near the ground surface (Figure 5). It is usually associated with intermediate or regional groundwater flow systems. If the pressure is high enough, the water flows to the surface and produces a flowing well, spring or soap hole. Artesian seeps can be identified from the presence of these flow features and from hydrogeological maps. Only one seep of this type occurs in the M.D. and it occupies less than 0.1% (2.4 ha) of the total saline area in the M.D. This seep is located in Twp. 28 Rge. 25, on the border with the County of Wheatland.




Figure 5. Artesian salinity

4. Depression Bottom Salinity

This salinity type occurs in low lying areas. Surface water is trapped temporarily in these areas until the water drains off and/or infiltrates the soil. Some water in the soil flows upslope through the upper soil in an unsaturated state and then surfaces to evaporate and deposit salt at the edge of the ponded area (Figure 6). As well, once the surface water has disappeared, groundwater from the water table rises by capillary action to the surface in and around the previously ponded area. Depression bottom seeps are well defined with distinct, rounded edges. In the M.D., this is the most common salinity type. It occupies 58.1% of the total saline area. Depression bottom salinity occurs in several landscapes with poor drainage, particularly in gently rolling areas. It is also associated with Solonetzic soils which occur throughout the M.D. The main areas of depression bottom salinity are: in the vicinity of Sheppard and Langdon (Twp. 23, 24 Rge. 27, 28 W4); and east of Irricana (Twp. 27 Rge. 25, 26 W4).




Figure 6. Depression bottom salinity

5. Coulee Bottom Salinity

Coulee bottom salinity forms in the bottoms of coulees and watercourses by the same mechanism as depression bottom salinity but on a larger scale. In the M.D. of Rocky View, it occupies only 0.9% of the total saline area. It occurs mainly along coulees including the MacPherson coulee and its tributaries (north of Airdrie), in the Beiseker area (Twp. 28 Rge. 25 W4), and near Crossfield Creek (Twp. 28 Rge. 28 W4). This salinity type typically develops over long periods of time so most of the lands affected by coulee bottom salinity have never been in agricultural production.

6. Slough Ring Salinity

This salinity type occurs as a ring of salt immediately adjacent to a permanent water body (Figure 7). Water infiltrates from the water body into the permeable upper soil layer and flows upslope through this layer as shallow groundwater in an unsaturated state. The water may also flow downward, raising the water table. Water from the unsaturated flow and water raised from the water table by capillary action emerges at the surface where it evaporates, leaving salts at the edge of the slough. Only three slough ring seeps occur in the M.D. Two are in Twp. 24 Rge. 28 W4 and one is in Twp. 27 Rge. 26 W4.




Figure 7. Slough ring salinity

2.2.2 Irrigation Salinity Types

1. Canal Seepage Salinity

This type of salinity is dominant in irrigated areas where leakage from canals contributes to seeps (Figure 8). Because many canals are located along a topographic break, canal seepage often aggravates natural salinity. This salinity type is the third most common type in the M.D., and occupies 15.7% of the total saline area. It occurs extensively in the Western Irrigation District, especially in the area southeast of Irricana (Twp. 26, 27 Rge. 25, 26 W4).




Figure 8. Canal seepage salinity

2. Natural/Irrigation Salinity

These seeps result from one or more of: natural seepage, canal seepage and excess irrigation. All seeps located on irrigated land and some distance from canals and supply ditches are given this classification. This salinity type occupies only 2.1% of the total saline area in the M.D. Only two seeps of this type occur in the M.D. and both are east of Irricana (Twp. 27 Rge. 25 W4).

2.3 Salinity Controls

Salinity is a complex problem caused by climatic, hydrogeological and agricultural factors. The opportunities for moderating the effects of climate and hydrogeological processes are limited and/or expensive. Therefore appropriate agricultural practices are used to help prevent or control saline seeps. The emphasis in this report is on cost-effective, agronomic measures. Specifically, cropping systems that intercept soil water in a recharge area before the water moves below the crop root zone are recommended.

Recommended control methods for the types of salinity found in the M.D. of Rocky View are summarized in Table 1 and described in more detail in Sections 2.3.1 and 2.3.2.

2.3.1 Biological Controls

Salt-Tolerant Crops

Saline areas should not be left bare for extended periods. Very saline soils should be seeded to a mixture of salt-tolerant forage crops (Table 2). Saline areas are often wet, so crops may need to be tolerant of both salt and excess moisture. When electrical conductivity measurements exceed 8 to 10 dS/m, salt-tolerant seed mixtures usually give the best results.

Establishing deep-rooted vegetation on a saline seep can be very difficult. Salt-tolerant grasses can be seeded in the fall when the saline seeps are dry and accessible. Seeding rates for saline seeps, especially when planted in the fall, should be double those for non-saline areas.

Table 1. Salinity types and control methods

Salinity type	Control
Contact/slope change	- salt-tolerant grasses in saline area and alfalfa in upslope recharge area (recharge area may be about three times area of seep)
Outcrop	- salt-tolerant grasses in saline area
Artesian	- salt-tolerant grasses in saline area - where applicable, install relief wells connected to suitable outlet
Depression bottom	- salt-tolerant grasses in saline area and along edge of depression in band 50 to 150 m (165 to 490 ft) wide - appropriate structural controls
Coulee bottom	- salt-tolerant grasses in saline area - appropriate structural controls
Slough ring	- deep-rooted and salt-tolerant grasses in a 20 to 60 m (65 to 195 ft) band around slough - appropriate structural controls
Irrigation canal seepage	- structural controls to prevent canal seepage (canal lining, cut-off curtains) and/or subsurface drainage of affected area
Natural/irrigation	- appropriate structural controls for irrigation-related salinity - salt-tolerant grasses for natural salinity

Deep-Rooted Crops

Deep-rooted crops prevent the buildup of groundwater, lower the water table, dry out the subsoil and restore the water storage capacity of the soil (Brown et al. 1982). The most commonly grown deep-rooted dryland crop is alfalfa. It roots up to 6 m (20 ft) in four to five years and uses more than 760 mm (30 in.) of water per year. Perennial deep-rooted crops also increase soil organic matter content, improve soil structure and reduce soil erosion.

Alfalfa should be seeded into a firm, moist seed bed as shallowly as possible at a rate of about 7 kg/ha (6.2 lbs/ac). It can be seeded using a conventional seeder. Hoe drills often give the most effective results because of good depth control and packing capability. Disk drills work best if the seed bed is uniform and moderately firm. However, in loose soil, disk drills may place the seed too deeply, and in very firm soil, they may leave the seed on the soil surface. Both conditions result in poor germination.

Recharge areas identified during a salinity investigation should be seeded to alfalfa. On average, recharge areas are about three times larger than their saline seep. The best time for seeding alfalfa is early spring. Alfalfa should be seeded without any companion crops because competition will deter establishment of the alfalfa stand.

Table 2. Forage crops for saline soils and flooded areas (Henry et al. 1987)

Salinity rating (EC)*	Forage mix	Seeding rate for hay or pasture (kg/ha)
a. Soils with little or no spring flooding (up to 2 weeks)		
Slight to moderate (2-6 dS/m)	bromegrass + Russian wild ryegrass + alfalfa (Rambler)	4+4+4
	bromegrass + slender wheatgrass + alfalfa (Rambler)	4+4+4
	Russian wild ryegrass + alfalfa	6+3
	altai wild ryegrass + alfalfa	10+3
	crested wheatgrass + alfalfa	7+3
	altai wild ryegrass	11
	slender wheatgrass + sweet clover (short-term stands and not over 1 week of flooding)	8+6
Severe (6-10 dS/m)	bromegrass + Russian wild ryegrass + slender wheatgrass	4+4+4
	altai wild ryegrass + alfalfa	10+3
	altai wild ryegrass	11
	tall wheatgrass (moist districts or seepage areas)	12
Very severe (10-15 dS/m)	Russian wild ryegrass + slender wheatgrass	4+4
	altai wild ryegrass + alfalfa	10+3
	altai wild ryegrass	10
	tall wheatgrass (moist districts or seepage areas)	12
b. Soils with spring flooding (2 to 5 weeks)		
Little or no (0-2 dS/m)	reed canarygrass + bromegrass	4+6
	reed canarygrass + timothy	4+4
	timothy + bromegrass	4+6
	altai wild ryegrass + alfalfa	10+3
	altai wild ryegrass	11
Slight to moderate (2-6 dS/m)	reed canarygrass + bromegrass	4+6
	reed canarygrass + bromegrass + slender wheatgrass	4+6+6
	altai wild ryegrass + alfalfa	10+3
	altai wild ryegrass	11
Severe to very severe (6-15 dS/m)	altai wild ryegrass + alfalfa	10+3
	slender wheatgrass	8
	altai wild ryegrass	11
	tall wheatgrass	12

* EC - electrical conductivity based on saturated paste, in deciSiemens per metre

Alfalfa has the ability to use atmospheric nitrogen through a symbiotic relationship with rhizobia nodule bacteria. Therefore, alfalfa inoculated with rhizobia will require less nitrogen fertilizer. Phosphorus, potassium and sulphur are important nutrients for optimum production. Alfalfa requires 5 kg of phosphorus per tonne of yield (10 lbs per ton of yield). This nutrient is very immobile in the soil and so application prior to seeding is highly recommended.

When alfalfa is seeded in a recharge area, it usually takes about five years to lower the water table in the associated saline seep. Once the water table is lowered to an acceptable level, the recharge area may be converted to cereal crops for a few years. The best approach is usually to establish a rotation of five years of alfalfa followed by three years of cereal crops. The cereals should be continuously cropped.

Flexible Cropping

In flexible cropping, fields are seeded if stored soil moisture and rainfall probabilities are favourable for satisfactory crop yields, and they are fallowed only if yield prospects are unfavourable (Jackson and Krall 1978). Flexible cropping involves careful management and planning; it is often simpler to use continuous cropping.

Snow trapping may increase stored soil moisture for recropping. Techniques to trap and manage snow include:

- tall stubble/alternate height stubble - Leaving tall stubble or strips of stubble at different heights increases stored soil moisture.
- shelterbelts - The ability of shelterbelts to trap snow can be manipulated by such practices as tree pruning and species selection.

Snow trapping should distribute snow evenly to avoid local accumulations of snowmelt from large drifts.

2.3.2 Structural Controls

Surface Drainage

Surface drainage of recharge areas and/or discharge areas can be used to control seeps (Vanderpluym 1982). An open, shallow trench is normally used; deep trenches will obstruct farming operations. Trenches can be constructed at a reasonable cost with farm equipment or by a contractor.

Grassed waterways are often used to drain excess surface water from recharge areas. Typical grassed waterways are broad, shallow channels with shallow slopes that carry water at slow speeds, preventing soil erosion. Grassed waterways ideally have channel slopes of less than 1% and side slopes of less than 25%. The channel should be at least 15 cm (5.9 in.) deep and 5 m (16 ft) wide. The grass should extend at least 5 m (16 ft) on both sides of the channel.

A commonly used forage mix for grassed waterways is:

- brome or pubescent wheatgrass - 10 kg/ha (9 lbs/ac), plus
- creeping red fescue - 5 kg/ha (4.5 lbs/ac), plus
- crested wheatgrass - 5 kg/ha (4.5 lbs/ac), plus
- alfalfa - 10 kg/ha (9 lbs/ac), plus
- fall rye - 1 kg/ha (0.9 lbs/ac)

Subsurface Drainage

Although subsurface (tile) drainage is used on irrigated lands to control salinity and waterlogging, it is not commonly used to control dryland salinity. However, a subsurface drainage system will satisfactorily lower water tables in dryland seeps if the system is properly designed, installed and managed. If the water is of good quality, it could be stored and used for stock water.

Relief Wells

Relief wells are costly but they can effectively control springs and soapholes associated with artesian salinity. The wells should be completed in the pressurized water-bearing layer. The wells may flow free and could be connected to a buried pipe 1 to 2 m (3.3 to 6.6 ft) deep. If the water is of good quality, it could be used for domestic or livestock purposes.

3.0 Saline Water Bodies


Salt concentration is an important water quality measure for cattle health and productivity. As part of the salinity mapping project, water bodies were sampled for salinity concentrations.

Livestock will normally avoid saline water if given the choice, but in some situations they may have to drink it. Use of water with total dissolved solids (TDS) concentrations higher than 3 000 mg/L should be restricted depending on livestock type, age and reproductive state (Task Force on Water Quality Guidelines 1987). Water with 7 000 mg/L TDS or more can cause a loss in overall condition and a reduction in weight gain for all livestock. Excessively saline water (greater than 10 000 mg/L), especially when used in large amounts, can cause physiological upset and ultimately death in the majority of terrestrial animals, including humans.


In most parts of Alberta, natural surface waters usually have TDS concentrations well below 3 000 mg/L. However, water bodies in some locations have TDS concentrations of 7 000 mg/L or more, high enough to be dangerous for cattle (MacAlpine and Howard 1995). If water with 7 000 to 10 000 mg/L TDS must be used, then its use should be limited to a short period of time. A sudden change from non-saline water to very saline water can initiate toxic symptoms and in some cases may cause death.

An EC reading of 8 dS/m is equivalent to 7 000 mg/L TDS. Thus, water bodies with greater than 8 dS/m are a potential danger for cattle. In the M.D. of Rocky View, electrical conductivity was measured in one permanent water body. At the time of the survey, all other water bodies in the dryland portion of the M.D. were dry. (Water bodies in the irrigated area had water but were not measured.) The sampled water body (shown on Figure 9) was measured at 43 dS/m and should be avoided for watering cattle. It should be noted that salinity and TDS concentrations will increase over the summer period if drought conditions persist.

**Figure 9. Saline water bodies and associated salinity types,
Twp. 27 Rge. 25 W4, M.D. of Rocky View**



☒ Saline water bodies showing electrical conductivity (EC) in dS/m



4.0 Saline Seep Distribution

The following statistical analysis describes the number and size of saline seeps for all types of salinity and for contact/slope change salinity in the M.D. of Rocky View. These two examples indicate the general tendencies for the other salinity types.

The M.D. has 915 saline seeps which occupy a total of 11 359 ha (28 068 ac). This project depicts salinity which is visible on the soil surface. Most of these visible saline areas are out of agricultural production or have significantly reduced crop yields. However, the effects of salinity on crop yields are not usually limited to the visible saline areas. Often the surrounding lands have weakly to very weakly saline subsoils, reducing yields of sensitive crops. Thus, salinity control practices may benefit crop yields over a much broader area than just the visible seep.

Table 3 presents the number, area and percentage of total saline land for each salinity type. Depression bottom salinity is the most common type (58.1% of saline land in the M.D.), followed by contact/slope change salinity (22.3% of saline land), irrigation canal seepage salinity (15.7%), and irrigation/natural (2.1%). Coulee bottom salinity occupies only 0.9% of the total saline land in the M.D. Outcrop salinity, slough ring salinity and artesian salinity are very minor types. Combined, they occupy only 0.9% of the saline land.

Table 3. Salinity distribution by type in the M.D. of Rocky View

Salinity type	Number of Seeps	Area		Percent of total saline area
		(ac)	(ha)	
Contact/slope change	554	6 257.3	2 532.2	22.3
Outcrop	13	168.1	68.0	0.6
Artesian	1	5.9	2.4	0.1
Depression bottom	263	16 311.2	6 600.9	58.1
Coulee bottom	8	258.1	104.5	0.9
Slough ring	3	65.8	26.6	0.2
Irrigation canal seepage	71	4 408.2	1 783.9	15.7
Natural/irrigation	2	593.7	240.3	2.1
Total	915	28 068.3	11 358.8	100.0

Figure 10 shows the number of saline seeps by type. Overall, there are many small contact/slope change saline seeps (554 seeps occupying 2 532 ha (6 257 ac)); there are fewer but larger depression bottom seeps (only 263 seeps occupying 6 600 ha or 16 311 ac). Irrigation canal seepage salinity consists of 71 saline seeps occupying 1 784 ha (4 408 ac). Outcrop salinity consists of 13 small seeps occupying 68 ha (168 ac). Coulee bottom salinity consists of 8 seeps occupying 104 ha (258 ac). There are only 3 slough ring seeps, 2 natural/irrigation seeps and 1 artesian seep.

Figure 11 shows the frequency of different size ranges for all saline seeps. The average seep size is 12.4 ha. The smallest seep is a contact/slope change seep with an area of 0.08 ha (0.19 ac.) and the largest is a depression bottom seep with an area of 679 ha (1 677 ac). Most seeps (52%) are between 0 and 3.75 ha (0 to 9.2 ac); more than 98% are between 0 and 100 ha (247 ac). Only 18 saline seeps are larger than 100 ha. Most seeps larger than 10 ha are depression bottom or coulee bottom seeps.

The areas for the contact/slope change saline seeps show a similar distribution (Figure 12). Fifty-seven percent are between 0 and 2.5 ha (0 and 6.2 ac), and 26.5% are greater than 4.5 ha (11.1 ac). The usual measures to control contact/slope change salinity are to grow salt-tolerant grasses in the saline area and alfalfa in the recharge area. On average, recharge areas are about three times the size of their saline seep. Thus, as a general guide, a recharge area about three times the size of the seep will need to be converted to alfalfa to control contact/slope change seeps.

The other salinity types have similar distributions of seep areas. In general, most saline seeps are between 0 and 3.75 ha. The main exceptions are some depression bottom and coulee bottom saline seeps.

Figure 10. Number of Saline Seeps by Type in the M.D. of Rockyview




Figure 11. Number and Size of Saline Seeps in the M.D. of Rockyview





Figure 12. Number and Size of Contact/Slope Change Saline Seeps in the M.D. of Rockyview



5.0 References

- Agriculture Canada. 1976. Glossary of terms in soil science. Agriculture Canada, Research Branch, Ottawa. Publication 1459, revised. 44 pp.
- Borneuf, D. 1972. Hydrogeology of the Drumheller area, Alberta. Research Council of Alberta. Report 72-1. 15 pp. 1 map.
- Brown, P.L., Halvorson, A.D., Siddoway, F.H., Mayland, H.F. and Miller, M.R. 1982. Saline seep diagnosis, control, and reclamation. United States Department of Agriculture. Conservation Research Report No. 30. 22 pp.
- Greenlee, G.M., Pawluk, S. and Bowser, W.E. 1968. Occurrence of soil salinity in the dry lands of southern Alberta. Canadian Journal of Soil Science. 48: 65-75.
- Henry, L., Harron, B., and Flaten, D. 1987. The nature and management of salt-affected land in Saskatchewan. Saskatchewan Agriculture. Agdex 518. 23 pp.
- Jackson, G.D. and Krall, J.L. 1978. The flexible method of recropping. Pages 7-23 *in* Dryland Saline Seep Control. Proceedings of the Meeting of the Subcommittee on Salt-Affected Soils at the 11th International Soil Science Society Congress. Edmonton.
- Kwiatkowski, J., Marciak, L.C, Wentz, D., and Vanderpluym, H. 1994. Salinity mapping by landscape, County of Vulcan, Alberta. Alberta Agriculture, Food and Rural Development and Agriculture and Agri-Food Canada, Edmonton.
- MacAlpine, N.D. and Howard, A. 1995. Farm water management manual. Conservation Manual Series. Alberta Agriculture, Food and Rural Development, Conservation and Development Branch, Edmonton.
- Ozoray, G.F. and Lytviak, A.T. 1974. Hydrogeology of the Gleichen area, Alberta. Alberta Research Council. Report 74-9. 16 pp. 1 map.
- Ozoray, G.F. and Barnes, R. 1978. Hydrogeology of the Calgary-Golden area, Alberta. Alberta Research Council. Report 77-2. 38 pp. 1 map.
- Prairie Farm Rehabilitation Administration. various dates. Salinity investigation reports. These reports cover 18 sites in the M.D. of Rocky View and were prepared in the 1980s and 1990s.
- Task Force on Water Quality Guidelines of the Canadian Council of Ministers of the Environment. 1987. Chapter 4.0 Agricultural Uses. *In*: Canadian Water Quality Guidelines. Environment Canada, Inland Waters Directorate, Water Quality Branch, Ottawa.
- Vanderpluym, H. 1982. Salinity in western Canada. Pages 9-23 *in* First Annual Western Provincial Conference of Water and Soil Research and Management, Soil Salinity. Warner.
- Vanderpluym, H. and Harron, B. 1992. Dryland salinity investigation procedures manual. Alberta Agriculture, Conservation and Development Branch, and Agriculture Canada, Prairie Farm Rehabilitation Administration.
- Western Irrigation District. 1990. Saline/waterlogged lands map. Alberta Agriculture, Irrigation and Resource Management Division, Land Evaluation and Reclamation Branch.

Wyatt, F.A., Newton, J.D., Bowser, W.E., and Odynsky, W. 1943. Soil Survey of Rosebud and Banff Sheets. University of Alberta. Bulletin 40. 126 pp. 2 maps.

Wyatt, F.A., Newton, J.D., Bowser, W.E., and Odynsky, W. 1960. Soil Survey Blackfoot and Calgary Sheets. University of Alberta. Bulletin SS-2 (revised edition of Bulletin No. 39), 44 pp. 2 maps.

Glossary

Aquifer - A body of earth material capable of transmitting water through its pores at a rate sufficient for water supply purposes. (Vanderpluym and Harron 1992)

Artesian groundwater - Groundwater confined under pressure so that water rises up in a non-pumping well which penetrates it. (Vanderpluym and Harron 1992)

Bedrock - The solid rock that underlies the soil and regolith or that is exposed at the surface. (Agriculture Canada 1976)

Capillary action - The action by which the surface of a liquid, where it is in contact with a solid, is elevated or depressed depending on the forces of adhesion and cohesion.

Electrical conductivity - A method of expressing salinity. An electrical conductivity (EC) measurement can be used to determine the salt content of soil in a saturated soil paste extract. The EC value is usually expressed in deciSiemens/metre (dS/m). For example, topsoil with an EC value of 2 dS/m is considered non-saline; topsoil with an EC value of 16 is very saline.

Flexible cropping - Cropping according to spring soil moisture conditions. That is, seeding when the spring soil moisture is adequate.

Groundwater - 1) Water that is passing through or standing in the soil and the underlying strata. It is free to move by gravity. (Agriculture Canada 1976). 2) Water in the ground that is in the zone of saturation, from which wells, springs and groundwater runoff are supplied. (Vanderpluym and Harron 1992)

Parent material - The unconsolidated and more or less chemically weathered mineral or organic matter from which the solum of a soil has developed. (Agriculture Canada 1976)

Permeability, soil - The ease with which gases and liquids penetrate or pass through a bulk mass of soil or a layer of soil. (Agriculture Canada 1976)

Saline soil - A non-sodic soil containing enough soluble salts to interfere with the growth of most crop plants. The conductivity of the saturation extract is greater than 4 dS/m (at 25°C), the exchangeable sodium percentage is less than 15, and the pH is usually less than 8.5. (Agriculture Canada 1976)

Seepage - 1) The emergence of water from the soil over an extensive area, in contrast to a spring where water emerges from a local spot. (Agriculture Canada 1976). 2) The slow movement of water through small cracks, pores, interstices, etc. of a material into or out of a body of surface or subsurface water (Vanderpluym and Harron 1992).

Till - Unstratified sediment deposited directly by a glacier and consisting of clay, sand, gravel and boulders intermingled in any proportion. (Agriculture Canada 1976)

For further information, please visit www.agric.gov.ab.ca